



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 7/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 00/02905</b> <b>(43) International Publication Date:</b> 20 January 2000 (20.01.00)
<b>(21) International Application Number:</b> PCT/US99/15735 <b>(22) International Filing Date:</b> 12 July 1999 (12.07.99)  <b>(30) Priority Data:</b> 60/092,653      13 July 1998 (13.07.98)      US 60/130,855      22 April 1999 (22.04.99)      US  <b>(71) Applicant (for all designated States except US):</b> UNIVERSITY OF SOUTHERN CALIFORNIA [US/US]; Suite 313, 3716 South Hope Street, Los Angeles, CA 90007-4344 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> RODGERS, Kathleen [US/US]; 4403 Galeano Street, Long Beach, CA 90815 (US). DIZEREGA, Gere [US/US]; 1270 Hillcrest Avenue, Pasadena, CA 91106 (US).  <b>(74) Agent:</b> HARPER, David, S.; McDonnell Boehnen Hulbert & Berghoff, 300 South Wacker Drive, Chicago, IL 60606 (US).		<b>(81) Designated States:</b> AU, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> METHODS FOR ACCELERATING BONE AND CARTILAGE GROWTH AND REPAIR  <b>(57) Abstract</b>  <p>The present invention provides improved methods, kits, and compositions for enhancing bone, cartilage and cartilage repair, bone and prosthesis implantation, and attachment and fixation of cartilage and cartilage to bone or other tissues, and chondrocyte proliferation comprising the administration of an effective amount of angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, angiotensin II (AII), AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

## METHODS FOR ACCELERATING BONE AND CARTILAGE GROWTH AND REPAIR

### 5 Cross Reference

This application is a continuation-in-part of U.S. Provisional Application Serial No. 60/092,653 filed July 13, 1998 and a continuation in part of U.S. Provisional Application Serial No. 60/130,855 filed March 8, 1999.

### 10 Field of the Invention

The present invention relates to methods, compositions, and kits for the repair, regeneration, and implantation of bone and cartilage.

### Background of the Invention

15 Natural mechanisms of repair, healing and augmentation are similar for bone and cartilage. (U.S. Patent No. 5,686,116) Although repair, healing and augmentation require a complex series of events that are not well defined, it is known that specific, naturally occurring factors are required to achieve these objectives. Such factors are released or migrate into the injured area, and stimulate osteoblasts and chondrocytes  
20 and odontoblasts in bone and cartilage to stimulate matrix formation and remodeling of the wounded area. (ten Dijke et al., Bio/Technology, 7:793-798 (1989))

Living bone tissue is continuously being replenished by the processes of resorption and deposition of bone matrix and minerals. This temporally and spatially coupled process, termed bone remodeling, is accomplished largely by two cell  
25 populations, the osteoclasts and osteoblasts. (U.S. Patent No. 5,656,598, incorporated

by reference herein in its entirety) The remodeling process is initiated when osteoclasts are recruited from the bone marrow or the circulation to the bone surface and remove a disk-shaped packet of bone. The bone matrix and mineral is subsequently replaced by a team of osteoblasts recruited to the resorbed bone surface from the bone marrow.

5 Osteoblasts are derived from local mesenchymal (stromal) precursors which differentiate into osteoblasts.

New bone can be formed by three basic mechanisms: osteogenesis, osteoconduction and osteoinduction. (U.S. Patent No. 5,464,439 incorporated by reference herein in its entirety) In osteogenic transplantation, viable osteoblasts and

10 peri-osteoblasts are moved from one body location to another where they establish centers of bone formation. Cancellous bone and marrow grafts provide such viable cells. TGF-beta has been shown to stimulate proliferation and matrix synthesis of osteoblastic cells (Centrella, et al. (1987) J. Biol. Chem. 262:2869-2874) and to inhibit the formation and activity of osteoclastic cells (Chenu, et al. (1988) Proc. Natl. Acad.

15 Sci. U.S.A. 85:683-5687; Kiebzak et al. (1988) J. Bone Min. Res. 3:439-446), and to stimulate local bone formation in vivo. (Joyce, et al. (1990) J. Cell. Biol. 110:2195-2207; Noda and Camilliere (1989) Endocrinology 124:2991-2294). Other factors reported to stimulate bone growth include bone morphogenetic proteins (WO 88/00205), insulin-like growth factor (IGF) (Endocrinol. Metab. 13:E367-72,1986), and

20 parathyroid hormone (J. Bone & Min. Res. 1:377-381, 1986).

Members of the bone morphogenetic protein family have been shown to be useful for induction of cartilage and bone formation. For example, BMP-2 has been shown to be able to induce the formation of new cartilage and/or bone tissue in vivo in a rat ectopic implant model, see U.S. Pat. No. 5,013,649; in mandibular defects in dogs,

see Toriumi et al., Arch. Otolaryngol Head Neck Surg., 117:1101-1112 (1991); and in femoral segmental defects in sheep, see Gerhart et al., Trans Orthop Res Soc, 16:172 (1991). Other members of the BMP family have also been shown to have osteogenic activity, including BMP-4, -6 and -7 (see Wozney, Bone Morphogenetic Proteins and Their Gene Expression, in Cellular and Molecular Biology of Bone, pp. 131-167 (Academic Press, Inc. 1993)). BMP proteins have also been shown to demonstrate inductive and/or differentiation potentiating activity on a variety of other tissues, including cartilage. (U.S. Patent No. 5,700,774, hereby incorporated by reference in its entirety.

10 In the transplantation of large segments of cortical bone or allogenic banked bone, direct osteogenesis does not occur. Rather, osteoconduction occurs wherein the dead bone acts as a scaffold for the ingrowth of blood vessels, followed by the resorption of the implant and deposition of new bone. This process is very slow however, often requiring years to reunite a large segmental defect.

15 Osteoinduction is the phenotypic conversion of connective tissue into bone by an appropriate stimulus. As this concept implies, formation of bone can be induced at even non-skeletal sites. Osteoinduction is preferred over osteoconduction, as grafts of this type are typically incorporated into the host bone within a two-week period. In contrast, osteoconductive grafts have been found to be non-incorporated as long as one  
20 year after implantation. In order to provide an environment suitable for osteoinduction, a material should be selected which is not only capable of inducing osteogenesis throughout its volume, but is also biocompatible, non-inflammatory, and possesses the ability to be ultimately resorbed by the body and replaced with new, natural bone.

Among the pathological conditions associated with abnormal bone cell function are osteoporosis, osteoarthritis, Paget's disease, osteomalacia, periodontal disease, bone loss resulting from multiple myeloma and other forms of cancer, bone loss resulting from side effects of other medical treatment (such as  
5 steroids), and age-related loss of bone mass. Inadequate organic matrix mass places an individual at risk of skeletal failure such that bone fractures can result from the minimal trauma of everyday life. Such fractures cause significant illness, or morbidity, inasmuch as there is insufficient repair or healing of the fractures. In certain pathologic conditions, osteoclast-mediated resorption is not regulated by osteoblasts but is driven  
10 by cancer cells, infecting organisms or the host's immune cells. In those disease conditions, resorption of bone far exceeds bone formation. Such accelerated osteoclastic activity leads to excessive release of calcium from the inorganic mineral in bone, with a concomitant net loss of skeletal mass, often with an attendant disturbance in calcium homeostasis in the form of elevated blood levels of calcium. (U.S. Patent  
15 No. 5,686,116, incorporated by reference herein in its entirety.)

Although methods for directing new bone formation are known, improved methods that provide for accelerated bone growth are needed. For example, currently approved therapeutic agents for osteoporosis are antiresorptives. As such, they are not as effective in patients with established osteoporosis of either type (decreased bone  
20 density with fractures of the vertebrae and/or hip), or in patients with Type II osteoporosis. In addition, the most accepted preventive agent for osteoporosis currently in use is estrogen therapy, which is not an acceptable therapeutic agent for women with a history of breast cancer or endometrial cancer or for men with osteoporosis.

Similarly, successful implantation and function of bone implants depends on bonding of the adjacent bone to the implant. (U.S. Patent No. 5,686,116) Such bonding requires bone repair by the formation of new matrix components at the interface between the implant and the bone proximate to the implant. An estimated ten percent  
5 of bone and joint prosthetic devices that are placed in people fail to function due to non-bonding of the bone to an implant. The resulting disability often requires reoperation and reimplantation of the device. Furthermore, five to ten percent of all bone fractures are never repaired. Although many methods have been proposed to cure these non-healing bone fractures, none has yet proven to be satisfactory. Based on all of  
10 the above, there clearly exists a need in the art for improved methods that provide for accelerated bone growth.

Cartilage is a specialized type of dense connective tissue consisting of cells embedded in a matrix. There are several kinds of cartilage. (U.S. Patent No. 5,736,372, herein incorporated by reference in its entirety.) Translucent cartilage having a  
15 homogeneous matrix containing collagenous fibers is found in articular cartilage, in costal cartilages, in the septum of the nose, in larynx and trachea. Articular cartilage is hyaline cartilage covering the articular surfaces of bones. Costal cartilage connects the true ribs and the sternum. Fibrous cartilage contains collagen fibers. Yellow cartilage is a network of elastic fibers holding cartilage cells which is primarily found in the  
20 epiglottis, the external ear, and the auditory tube. Cartilage is tissue made up of extracellular matrix primarily comprised of the organic compounds collagen, hyaluronic acid (a proteoglycan), and chondrocyte cells, which are responsible for cartilage production. Collagen, hyaluronic acid and water entrapped within these organic matrix elements yield the unique elastic properties and strength of cartilage.

Chondrocytes produce both Type I and Type II collagens. Type II collagen is not found in bone, whereas Type I collagen is found in bone. (U.S. Patent No. 5,686,116) It has previously been shown that the endogenous growth factors TGF beta and BMP induce both new cartilage and bone formation. Wozney et al. Science, 242:1528-1533 (1988) and Sporn et al. J. Cell Biol. 105:1039-1045 (1987).

In cartilage, collagen synthesis is required for repair, healing and augmentation, as well as for the successful bonding of grafts and prosthetic devices. (U.S. Patent No. 5,686,116) Collagen is the major structural protein responsible for the architectural integrity of cartilage. Thus, an adequate supply of chondrocytes is essential in order to produce sufficient amounts of collagen for repair, healing, and augmentation of cartilage. Other, noncollagen proteins, such as osteonectin, fibronectin and proteoglycans are also important for cartilage repair.

Cells such as synoviocytes that are found in joint spaces adjacent to cartilage have an important role in cartilage metabolism. Synoviocytes produce metallo-proteinases, such as collagenases that are capable of breaking-down cartilage. TGF beta is known to inhibit cell-release (and probably synthesis) of metallo-proteinases and to induce chondrocytes (cartilage forming cells) to produce new matrix components and inhibit production of cartilage destructive enzymes so as to effect cartilage repair, healing and augmentation. Sporn et al. (1987). It has also been shown that mice deficient in parathyroid hormone-related peptide (PTHrP) exhibit abnormal cartilage maturation, indicating that PTHrP is an essential factor for chondrocyte development and maturation. (U.S. Patent No. 5,700,774)

Cartilage implants are often used in reconstructive or plastic surgery such as rhinoplasty. There is a need in the art for methods that increase chondrocyte



proliferation and collagen synthesis, and thus inhibit cartilage destruction and enhance cartilage repair. Such methods would increase the clinical utility of cartilage repair including but not limited to cartilage grafts and healing of cartilage grafts.

Although some of the above methods have met with limited success, there  
5 remains a need in the art for improved methods for enhancing bone and cartilage repair, healing and augmentation, and for enhancing the attachment and fixation of bone and cartilage implants.

### Summary of the Invention

10 The present invention provides methods, kits, and compositions for 1) enhancing bone and cartilage repair; 2) bone and prosthesis implantation; 3) attachment and fixation of cartilage to bone or other tissues; and methods, cell culture medium and kits for the proliferation of chondrocytes; all of which comprise the administration of  
15 angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, angiotensin II (AII), AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists.

These aspects and other aspects of the invention become apparent in light of the following detailed description.

### Brief Description of the Figures

Figure 1 is a bar graph showing the effect of AII, AIII, and GSD37B (10 µg/ml) on chondrocyte proliferation.

Figure 2 is a bar graph showing the effect of AII, GSD36, GSD37B, GSD38B, and  
25 GSD28 (10 µg/ml) of the invention on chondrocyte proliferation.

Figure 3 is a bar graph showing the effect of AII, 1GD, 2GD, and 3GD (10µg/ml) on chondrocyte proliferation.

Figure 4 is a bar graph showing the effect of AII, AII(1-7), GSD22A, GSD24B, and GSD28 (10 µg/ml) of the invention on chondrocyte proliferation.

5

### Detailed Description of the Preferred Embodiments

All references, patents and patent applications are hereby incorporated by reference in their entirety.

Within this application, unless otherwise stated, the techniques utilized may be found in any of several well-known references such as: *Molecular Cloning: A Laboratory Manual* (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press), *Gene Expression Technology* (Methods in Enzymology, Vol. 185, edited by D. Goeddel, 1991. Academic Press, San Diego, CA), "Guide to Protein Purification" in *Methods in Enzymology* (M.P. Deutscher, ed., (1990) Academic Press, Inc.); *PCR Protocols: A Guide to Methods and Applications* (Innis, et al. 1990. Academic Press, San Diego, CA), *Culture of Animal Cells: A Manual of Basic Technique, 2<sup>nd</sup> Ed.* (R.I. Freshney. 1987. Liss, Inc. New York, NY), *Gene Transfer and Expression Protocols*, pp. 109-128, ed. E.J. Murray, The Humana Press Inc., Clifton, N.J.), and the Ambion 1998 Catalog (Ambion, Austin, TX).

As defined herein the phrase "enhancing bone repair" refers to increasing the rate of new bone formation via bone remodeling, osteogenesis, osteoconduction and/or osteoinduction. The methods for enhancing bone repair in a mammal of the invention include those that stimulate bone formation and those that reverse bone loss. The methods can thus be used for (1) providing a subject with an amount of a substance

sufficient to act prophylactically to prevent the development of a weakened and/or unhealthy state; or (2) providing a subject with a sufficient amount of a substance so as to alleviate or eliminate a disease state and/or the symptoms of a disease state, and a weakened and/or unhealthy state.

5           As used herein the term "enhancing cartilage repair" comprises healing and regeneration of cartilage injuries, tears, deformities or defects, and prophylactic use in preventing damage to cartilaginous tissue.

          The present invention fulfills the need for methods to enhance bone repair in a mammal suffering from bone fractures, defects, and disorders which result in weakened  
10   bones such as osteoporosis, osteoarthritis, Paget's disease, osteomalacia, osteomalacia, periodontal disease, bone loss resulting from multiple myeloma and other forms of cancer, bone loss resulting from side effects of other medical treatment (such as steroids), and age-related loss of bone mass. In addition, bony ingrowth into various prosthetic devices can be greatly enhanced so that such artificial parts are firmly and  
15   permanently anchored into the surrounding skeletal tissue through a natural osseous bridge.

          The present invention further fulfills the need for methods to enhance the repair of cartilage in a mammal, by accelerating the proliferation of chondrocytes and thereby increasing the synthesis of collagen for use in cartilage repair. Such methods have  
20   application in the healing of cartilage, for example articular cartilage tears, deformities and other cartilage defects in humans and other animals. The methods have prophylactic use in preventing damage to cartilaginous tissue, as well as use in the improved fixation of cartilage to bone or other tissues, and in repairing defects to cartilage tissue. De novo cartilaginous tissue formation induced by the compounds of

the present invention contributes to the repair of congenital, trauma induced, or other cartilage defects of other origin, and is also useful in surgery for attachment or repair of cartilage. The methods and compositions of the invention may also be useful in the treatment of arthritis and other cartilage defects. The methods of the present invention  
5 can also be used in other indications wherein it is desirable to heal or regenerate cartilage tissue. Such indications include, without limitation, regeneration or repair of injuries to the articular cartilage. The methods of the present invention provide an environment to attract cartilage-forming cells, stimulate growth of cartilage-forming cells or induce differentiation of progenitors of cartilage-forming cells and  
10 chondrocytes.

The methods and kits of the present invention also provide improved chemically defined medium for accelerating the proliferation of chondrocytes (cartilage-forming cells). In another embodiment, the compositions and methods of the present invention can be used to treat chondrocytic cell lines, such as articular chondrocytes, in order to  
15 maintain chondrocytic phenotype and survival of the cells. The treated cell populations are therefore also useful for gene therapy applications.

U.S. Patent No. 5,015,629 to DiZerega (the entire disclosure of which is hereby incorporated by reference) describes a method for increasing the rate of healing of wound tissue, comprising the application to such tissue of angiotensin II (AII) in an  
20 amount which is sufficient for said increase. The application of AII to wound tissue significantly increases the rate of wound healing, leading to a more rapid re-epithelialization and tissue repair. The term AII refers to an octapeptide present in humans and other species having the sequence Asp-Arg-Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:1]. The biological formation of angiotensin is initiated by the action of

renin on the plasma substrate angiotensinogen (Clouston et al., *Genomics* 2:240-248 (1988); Kageyama et al., *Biochemistry* 23:3603-3609; Ohkubo et al., *Proc. Natl. Acad. Sci.* 80:2196-2200 (1983); each reference hereby incorporated in its entirety). The substance so formed is a decapeptide called angiotensin I (AI) which is converted to  
5 AII by the angiotensin converting enzyme (ACE) which removes the C-terminal His-Leu residues from AI [SEQ ID NO: 37]. AII is a known pressor agent and is commercially available.

Studies have shown that AII increases mitogenesis and chemotaxis in cultured cells that are involved in wound repair, and also increases their release of growth  
10 factors and extracellular matrices (diZerega, U.S. Patent No. 5,015,629; Dzau et. al., *J. Mol. Cell. Cardiol.* 21:S7 (Supp III) 1989; Berk et. al., *Hypertension* 13:305-14 (1989); Kawahara, et al., *BBRC* 150:52-9 (1988); Naftilan, et al., *J. Clin. Invest.* 83:1419-23 (1989); Taubman et al., *J. Biol. Chem.* 264:526-530 (1989); Nakahara, et al., *BBRC* 184:811-8 (1992); Stouffer and Owens, *Circ. Res.* 70:820 (1992); Wolf, et al., *Am. J.*  
15 *Pathol.* 140:95-107 (1992); Bell and Madri, *Am. J. Pathol.* 137:7-12 (1990). In addition, AII was shown to be angiogenic in rabbit corneal eye and chick chorioallantoic membrane models (Fernandez, et al., *J. Lab. Clin. Med.* 105:141 (1985); LeNoble, et al., *Eur. J. Pharmacol.* 195:305-6 (1991). Additionally, AII and angiotensin III analogs and fragments thereof have been shown to be effective in  
20 wound healing. (U.S. Patent No. 5,629,292; International Application No. WO 95/08565; International Application WO 95/08337; International Application No. WO 96/39164; all references hereby incorporated in their entirety.)

Previous studies have suggested that angiotensin I (AI) and AII both stimulate bone resorption *in vitro* by osteoclasts incubated on bone slices, but only in the

presence of osteoblastic cells, suggesting that the effect of angiotensin II was not direct, but rather is mediated by a primary hormonal interaction on cells of the osteoblastic lineage. (Hatton et al., *J. Endocrinol.* 152:5-10 (1997)). AI stimulation of bone resorption was inhibited by ACE inhibitors, suggesting that the formation of AII from AI was responsible for the stimulation of bone resorption. Neither AI nor AII were shown to have any effect on osteoclast formation. Thus, this study suggests that local bone destruction may be mediated by AII's stimulation of bone resorption.

Other studies have demonstrated AII stimulation of DNA and collagen synthesis *in vitro* on primary cultures of isolated, phenotypically immature osteoblasts derived from the periosteum of fetal rat calvariae and human adult trabecular bone. (Lamparter et al., *J. Cell. Physiol.* 175:89-98 (1998)) No direct AII effect was detected on primary cell populations with a mature osteoblast phenotype, and an indirect effect through AII-responsive osteoblastic precursor cells was proposed. Similar *in vitro* studies on osteoblast-rich populations of cells demonstrated a similar effect, while not ruling out stimulation of mature osteoblast proliferation. (Hiruma et al., *Biochem and Biophys. Res. Commun.* 230:176-178 (1997)) Another study suggests that AII may decelerate the differentiation and bone formation of rat calvarial osteoblasts. (Hagiwara et al., *J. of Endocrinology* 156:543-550 (1998))

Based on all of the above studies, there is no expectation that the use of angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, AII, AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists would be effective in enhancing bone and cartilage repair, or effective in accelerating chondrocyte proliferation and collagen synthesis.

Previous studies in our laboratory have demonstrated that a class of AII and AII analogues and fragments stimulate the proliferation of mesenchymal stem cells, which give rise to the cells that make up bone and cartilage. (U.S Patent Application 09/012,400, filed January 23, 1998, herein incorporated by reference in its entirety.)

5 A peptide agonist selective for the AT2 receptor (AII has 100 times higher affinity for AT2 than AT1) has been identified. This peptide is p-aminophenylalanine 6-AII ["(p-NH<sub>2</sub>-Phe)6-AII"], Asp-Arg-Val-Tyr-Ile-Xaa-Pro-Phe [SEQ ID NO.36] wherein Xaa is p-NH<sub>2</sub>-Phe (Speth and Kim, BBRC 169:997-1006 (1990). This peptide gave binding characteristics comparable to AT2 antagonists in the experimental models  
10 tested (Catalioto, et al., *Eur. J. Pharmacol.* 256:93-97 (1994); Bryson, et al., *Eur. J. Pharmacol.* 225:119-127 (1992).

The effects of AII receptor and AII receptor antagonists have been examined in two experimental models of vascular injury and repair which suggest that both AII receptor subtypes (AT1 and AT2) play a role in wound healing (Janiak et al.,  
15 *Hypertension* 20:737-45 (1992); Prescott, et al., *Am. J. Pathol.* 139:1291-1296 (1991); Kauffman, et al., *Life Sci.* 49:223-228 (1991); Viswanathan, et al., *Peptides* 13:783-786 (1992); Kimura, et al., *BBRC* 187:1083-1090 (1992).

Many studies have focused upon AII(1-7) (AII residues 1-7) or other fragments of AII to evaluate their activity. AII(1-7) elicits some, but not the full range of effects  
20 elicited by AII. Pfeilschifter, et al., *Eur. J. Pharmacol.* 225:57-62 (1992); Jaiswal, et al., *Hypertension* 19(Supp. II):II-49-II-55 (1992); Edwards and Stack, *J. Pharmacol. Exper. Ther.* 266:506-510 (1993); Jaiswal, et al., *J. Pharmacol. Exper. Ther.* 265:664-673 (1991); Jaiswal, et al., *Hypertension* 17:1115-1120 (1991); Portsi, et al., *Br. J. Pharmacol.* 111:652-654 (1994).

As hereinafter defined, a preferred class of AT<sub>2</sub> agonists for use in accordance with the present invention comprises angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, AII, AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists having p-NH-Phe in a position  
5 corresponding to a position 6 of AII. In addition to peptide agents, various nonpeptidic agents (e.g., peptidomimetics) having the requisite AT<sub>2</sub> agonist activity are further contemplated for use in accordance with the present invention.

The active AII analogues, fragments of AII and analogues thereof of particular interest in accordance with the present invention comprise a sequence consisting of at  
10 least three contiguous amino acids of groups R<sup>1</sup>-R<sup>8</sup> in the sequence of general formula I



in which R<sup>1</sup> and R<sup>2</sup> together form a group of formula



15 wherein X is H or a one to three peptide group, or is absent,

R<sup>A</sup> is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

R<sup>B</sup> is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;  
20

R<sup>3</sup> is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

R<sup>4</sup> is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;



R<sup>5</sup> is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

R<sup>6</sup> is His, Arg or 6-NH<sub>2</sub>-Phe;

R<sup>7</sup> is Pro or Ala; and

5 R<sup>8</sup> is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including R<sup>4</sup> as a terminal Tyr group, or is absent.

Compounds falling within the category of AT2 agonists useful in the practice of the invention include the AII analogues set forth above subject to the restriction that R<sup>6</sup> is p-NH<sub>2</sub>-Phe.

10 Particularly preferred combinations for R<sup>A</sup> and R<sup>B</sup> are Asp-Arg, Asp-Lys, Glu-Arg and Glu-Lys. Particularly preferred embodiments of this class include the following: AII, AIII or AII(2-8), Arg-Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:2]; AII(3-8), also known as des1-AIII or AIV, Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:3]; AII(1-7), Asp-Arg-Val-Tyr-Ile-His-Pro [SEQ ID NO:4]; AII(2-7), Arg-Val-Tyr-Ile-His-Pro [SEQ ID NO:5]; AII(3-7), Val-Tyr-Ile-His-Pro [SEQ ID NO:6]; AII(5-8), Ile-His-Pro-Phe [SEQ ID NO:7]; AII(1-6), Asp-Arg-Val-Tyr-Ile-His [SEQ ID NO:8]; AII(1-5), Asp-Arg-Val-Tyr-Ile [SEQ ID NO:9]; AII(1-4), Asp-Arg-Val-Tyr [SEQ ID NO:10]; and AII(1-3), Asp-Arg-Val [SEQ ID NO:11]. Other preferred embodiments include: Arg-norLeu-Tyr-Ile-His-Pro-Phe [SEQ ID NO:12] and Arg-Val-Tyr-norLeu-His-Pro-Phe [SEQ ID NO:13]. Still another preferred embodiment encompassed within the scope of the invention is a peptide having the sequence Asp-Arg-Pro-Tyr-Ile-His-Pro-Phe [SEQ ID NO:31]. AII(6-8), His-Pro-Phe [SEQ ID NO:14] and AII(4-8), Tyr-Ile-His-Pro-Phe [SEQ ID NO:15] were also tested and found not to be effective.

20

In a particularly preferred embodiment of the methods for chondrocyte proliferation, collagen synthesis, cartilage repair, and attachment and fixation of cartilage to bone or other tissues, the active compounds of the present invention are selected from those comprising the following general formula:

5 R1-R2-R3-R4-R5-His-Pro-R6, wherein

R1 is selected from the group consisting of Hydrogen, Gly, and Asp;

R2 is selected from the group consisting of Arg, Citron, or Ornithine;

R3 is selected from the group consisting of Val, Ile, Ala, Leu, and norLeu, or Pro;

10 R4 is selected from Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, and Ala;

R5 is selected from the group consisting of Ile, Ala, Val, Leu, and norLeu; and

R6 is Phe, Ile, or is absent.

Most particularly preferred embodiments of this class of compounds are selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 13, SEQ ID NO: 18, SEQ ID NO:19, SEQ ID NO:12, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:45.

20 In a particularly preferred embodiment of the methods for bone repair and bone and prosthesis implantation, the active compounds of the present invention are selected from those comprising the following general formula:

Asp-Arg-R1-R2-Ile-His-Pro-R2, wherein


R1 is selected from the group consisting of Ile, Pro, Ala, Val, Leu, and norLeu;

R2 is selected from Tyr and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R<sup>3</sup> is Phe, or is absent.

Most particularly preferred embodiments of this class of compounds are selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:24, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO: 33, SEQ ID NO:41, and SEQ ID NO:  
5 45.

Another class of compounds of particular interest in accordance with the present invention are those of the general formula II



in which R<sup>2</sup> is selected from the group consisting of H, Arg, Lys, Ala,  
10 Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

R<sup>3</sup> is selected from the group consisting of Val, Ala, Leu, norLeu, Ile,  
Gly, Pro, Aib, Acpc and Tyr;

R<sup>4</sup> is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ser,  
homoSer, Ala, and azaTyr;


15 R<sup>5</sup> is selected from the group consisting of Ile, Ala, Leu, norLeu, Val  
and Gly;

R<sup>6</sup> is His, Arg or 6-NH<sub>2</sub>-Phe;

R<sup>7</sup> is Pro or Ala; and

R<sup>8</sup> is selected from the group consisting of Phe, Phe(Br), Ile and Tyr.

20 A particularly preferred subclass of the compounds of general formula II has the  
formula



wherein R<sup>2</sup>, R<sup>3</sup> and R<sup>5</sup> are as previously defined. Particularly preferred is angiotensin III of the formula Arg-Val-Tyr-Ile-His-Pro-Phe [SEQ ID NO:2]. Other preferred compounds include peptides having the structures Arg-Val-Tyr-Gly-His-Pro-Phe [SEQ ID NO:17] and Arg-Val-Tyr-Ala-His-Pro-Phe [SEQ ID NO:18]. The  
 5 fragment AII(4-8) was ineffective in repeated tests; this is believed to be due to the exposed tyrosine on the N-terminus.

In the above formulas, the standard three-letter abbreviations for amino acid residues are employed. In the absence of an indication to the contrary, the L-form of the amino acid is intended. Other residues are abbreviated as follows:

10 TABLE 1

## Abbreviation for Amino Acids

Me <sup>2</sup> Gly	N,N-dimethylglycyl
Bet	1-carboxy-N,N,N-trimethylmethanaminium hydroxide inner salt (betaine)
Suc	Succinyl
Phe(Br)	p-bromo-L-phenylalanyl
azaTyr	aza- $\alpha'$ -homo-L-tyrosyl
Acpc	1-aminocyclopentane carboxylic acid
Aib	2-aminoisobutyric acid
Sar	N-methylglycyl (sarcosine)
Orn	Ornithine

It has been suggested that AII and its analogues adopt either a *gamma* or a *beta* turn (Regoli, et al., *Pharmacological Reviews* 26:69 (1974)). In general, it is believed that neutral side chains in position R<sup>3</sup>, R<sup>5</sup> and R<sup>7</sup> may be involved in maintaining the

appropriate distance between active groups in positions R<sup>4</sup>, R<sup>6</sup> and R<sup>8</sup> primarily responsible for binding to receptors and/or intrinsic activity. Hydrophobic side chains in positions R<sup>3</sup>, R<sup>5</sup> and R<sup>8</sup> may also play an important role in the whole conformation of the peptide and/or contribute to the formation of a hypothetical hydrophobic pocket.

5        Appropriate side chains on the amino acid in position R<sup>2</sup> may contribute to affinity of the compounds for target receptors and/or play an important role in the conformation of the peptide. For this reason, Arg and Lys are particularly preferred as R<sup>2</sup>.

For purposes of the present invention, it is believed that R<sup>3</sup> may be involved in the formation of linear or nonlinear hydrogen bonds with R<sup>5</sup> (in the gamma turn model) or R<sup>6</sup> (in the beta turn model). R<sup>3</sup> would also participate in the first turn in a beta antiparallel structure (which has also been proposed as a possible structure). In contrast to other positions in general formula I, it appears that beta and gamma branching are equally effective in this position. Moreover, a single hydrogen bond may be sufficient to maintain a relatively stable conformation. Accordingly, R<sup>3</sup> may suitably be selected from Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr.

10  
15

With respect to R<sup>4</sup>, conformational analyses have suggested that the side chain in this position (as well as in R<sup>3</sup> and R<sup>5</sup>) contribute to a hydrophobic cluster believed to be essential for occupation and stimulation of receptors. Thus, R<sup>4</sup> is preferably selected from Tyr, Thr, Tyr (PO<sub>3</sub>)<sub>2</sub>, homoSer, Ser and azaTyr. In this position, Tyr is particularly preferred as it may form a hydrogen bond with the receptor site capable of accepting a hydrogen from the phenolic hydroxyl (Regoli, et al. (1974), *supra*). R<sup>3</sup> may also be suitably Ala.

20

In position R<sup>5</sup>, an amino acid with a  $\beta$  aliphatic or alicyclic chain is particularly desirable. Therefore, while Gly is suitable in position R<sup>5</sup>, it is preferred that the amino acid in this position be selected from Ile, Ala, Leu, norLeu, Gly and Val.

In the angiotensinogen, AI, AI analogues, AI fragments and analogues thereof, AII analogues, fragments and analogues of fragments of particular interest in accordance with the present invention, R<sup>6</sup> is His, Arg or 6-NH<sub>2</sub>-Phe. The unique properties of the imidazole ring of histidine (e.g., ionization at physiological pH, ability to act as proton donor or acceptor, aromatic character) are believed to contribute to its particular utility as R<sup>6</sup>. For example, conformational models suggest that His may participate in hydrogen bond formation (in the *beta* model) or in the second turn of the antiparallel structure by influencing the orientation of R<sup>7</sup>. Similarly, it is presently considered that R<sup>7</sup> should be Pro in order to provide the most desirable orientation of R<sup>8</sup>. In position R<sup>8</sup>, both a hydrophobic ring and an anionic carboxyl terminal appear to be particularly useful in binding of the analogues of interest to receptors; therefore, Tyr and especially Phe are preferred for purposes of the present invention.

Analogues of particular interest include the following:

TABLE 2

## Angiotensin II Analogues

AII Analogue Name	Amino Acid Sequence	Sequence Identifier
Analogue 1	Asp-Arg-Val-Tyr-Val-His-Pro-Phe	SEQ ID NO: 19
Analogue 2	Asn-Arg-Val-Tyr-Val-His-Pro-Phe	SEQ ID NO: 20
Analogue 3	Ala-Pro-Gly-Asp-Arg-Ile-Tyr-Val-His-Pro-Phe	SEQ ID NO: 21
Analogue 4	Glu-Arg-Val-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 22
Analogue 5	Asp-Lys-Val-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 23
Analogue 6	Asp-Arg-Ala-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 24
Analogue 7	Asp-Arg-Val-Thr-Ile-His-Pro-Phe	SEQ ID NO: 25
Analogue 8	Asp-Arg-Val-Tyr-Leu-His-Pro-Phe	SEQ ID NO: 26

Analogue 9	Asp-Arg-Val-Tyr-Ile-Arg-Pro-Phe	SEQ ID NO: 27
Analogue 10	Asp-Arg-Val-Tyr-Ile-His-Ala-Phe	SEQ ID NO: 28
Analogue 11	Asp-Arg-Val-Tyr-Ile-His-Pro-Tyr	SEQ ID NO: 29
Analogue 12	Pro-Arg-Val-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 30
Analogue 13	Asp-Arg-Pro-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 31
Analogue 14	Asp-Arg-Val-Tyr(PO <sub>3</sub> ) <sub>2</sub> -Ile-His-Pro-Phe	SEQ ID NO: 32
Analogue 15	Asp-Arg-norLeu-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 33
Analogue 16	Asp-Arg-Val-Tyr-norLeu-His-Pro-Phe	SEQ ID NO: 34
Analogue 17	Asp-Arg-Val-homoSer-Tyr-Ile-His-Pro-Phe	SEQ ID NO: 35

The polypeptides of the instant invention may be synthesized by any conventional method, including, but not limited to, those set forth in J. M. Stewart and J. D. Young, *Solid Phase Peptide Synthesis*, 2nd ed., Pierce Chemical Co., Rockford, Ill. (1984) and J. Meienhofer, *Hormonal Proteins and Peptides*, Vol. 2, Academic Press, New York, (1973) for solid phase synthesis and E. Schroder and K. Lubke, *The Peptides*, Vol. 1, Academic Press, New York, (1965) for solution synthesis. The disclosures of the foregoing treatises are incorporated by reference herein.

In general, these methods involve the sequential addition of protected amino acids to a growing peptide chain (U.S. Patent No. 5,693,616, herein incorporated by reference in its entirety). Normally, either the amino or carboxyl group of the first amino acid and any reactive side chain group are protected. This protected amino acid is then either attached to an inert solid support, or utilized in solution, and the next amino acid in the sequence, also suitably protected, is added under conditions amenable to formation of the amide linkage. After all the desired amino acids have been linked in the proper sequence, protecting groups and any solid support are removed to afford the crude polypeptide. The polypeptide is desalted and purified, preferably chromatographically, to yield the final product.

Preferably, peptides are synthesized according to standard solid-phase methodologies, such as may be performed on an Applied Biosystems Model 430A

peptide synthesizer (Applied Biosystems, Foster City, Calif.), according to manufacturer's instructions. Other methods of synthesizing peptides or peptidomimetics, either by solid phase methodologies or in liquid phase, are well known to those skilled in the art.

5 In one aspect, the present invention provides methods and kits for enhancing bone and cartilage repair, implantation, and augmentation in a mammal comprising the administration of angiotensinogen, angiotensin I (AI), AI analogues, AI fragments and analogues thereof, angiotensin II (AII), AII analogues, AII fragments or analogues thereof or AII AT<sub>2</sub> type 2 receptor agonists (hereinafter referred to as "active agents").

10 The compounds can be administered alone or in combination with other compounds that enhance bone and/or cartilage repair, implantation, and augmentation, including but not limited to bone morphogenic protein-2, bone morphogenic protein-4, bone morphogenic protein-6, bone morphogenic protein-7, transforming growth factor-beta, insulin-like growth factor, and parathyroid hormone.

15 The active agents may be administered by any suitable route, including orally, parentally, by inhalation spray, rectally, or topically in dosage unit formulations containing conventional pharmaceutically acceptable carriers, adjuvants, and vehicles. Such vehicles may include a tantalum or hydroxyapatite scaffold as a vehicle with the compounds of the invention embedded therein. Alternatively, polymeric substrates can

20 be used for compound delivery to the bone or cartilage such as the polymeric substrates disclosed in U.S. Patent Nos. 5,443,515; 5,171,273; 5,607,474; 4,916,207; and 5,324,775; all references hereby incorporated in their entirety. The term parenteral as used herein includes topical (i.e.: placement in to the bone), subcutaneous, intravenous,



intraarterial, intramuscular, intrasternal, intratendinous, intraspinal, intracranial, intrathoracic, infusion techniques or intraperitoneally.

The active agents of the present invention can also be incorporated into a coating on the surface of a prosthetic device. Such coatings may be composed of a polymer that allows slow diffusion of the active agents at a rate sufficient to enhance bone attachment for a suitable period of time. Suitable coatings include, but are not limited to, hydroxyapatite, methacrylate and tricalcium phosphate. Further, the polymeric coatings can be applied only to the sites on the prosthetic device where bony ingrowth is desired. Bone grafts can be coated with or soaked or immersed in a rinse or gel prior to implantation so as to impregnate the graft with the active agents. Topical or local administration of the active agents to either the site of implantation or the implant itself, is preferred, as it diminishes drug exposure for tissues and organs not requiring treatment.

The active agents may be made up in a solid form (including granules, powders or suppositories) or in a liquid form (*e.g.*, solutions, suspensions, or emulsions). The compounds of the invention may be applied in a variety of solutions. Suitable solutions for use in accordance with the invention are sterile, dissolve sufficient amounts of the peptide, and are not harmful for the proposed application. In this regard, the compounds of the present invention are very stable but are hydrolyzed by strong acids and bases. The compounds of the present invention are soluble in organic solvents and in aqueous solutions at pH 5-8.

The active agents may be subjected to conventional pharmaceutical operations such as sterilization and/or may contain conventional adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers, buffers etc.

For administration, the active agents are ordinarily combined with one or more adjuvants appropriate for the indicated route of administration. The compounds may be admixed with lactose, sucrose, starch powder, cellulose esters of alcanoic acids, stearic acid, talc, magnesium stearate, magnesium oxide, sodium and calcium salts of phosphoric and sulphuric acids, acacia, gelatin, sodium alginate, polyvinylpyrrolidone, and/or polyvinyl alcohol, and tableted or encapsulated for conventional administration. Alternatively, the compounds of this invention may be dissolved in saline, water, polyethylene glycol, propylene glycol, carboxymethyl cellulose colloidal solutions, ethanol, corn oil, peanut oil, cottonseed oil, sesame oil, tragacanth gum, and/or various buffers. Other adjuvants and modes of administration are well known in the pharmaceutical art. The carrier or diluent may include time delay material, such as glyceryl monostearate or glyceryl distearate alone or with a wax, or other materials well known in the art.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin (e.g., liniments, lotions, ointments, creams, or pastes) and drops suitable for administration to the eye, ear, or nose.

The dosage regimen for enhancing bone and cartilage repair with the active agents is based on a variety of factors, including the type of injury, the age, weight, sex, medical condition of the individual, the severity of the condition, the route of administration, and the particular compound employed. Thus, the dosage regimen may vary widely, but can be determined routinely by a physician using standard methods. Dosage levels of the order of between 0.1 ng/kg and 10 mg/kg body weight of the active agents are useful for all methods of use disclosed herein.

As an example, the dosage regimen for accelerating bone and cartilage repair with the active agents wherein the compounds are embedded in a scaffold used for bone and/or cartilage, such as tantalum or hydroxyapatite, would be based upon the volume of bone to be filled and not upon the weight of the subject being treated. Following  
5 determination of the specific dose volume to be administered to the subject, calculated based on the defect, individual doses are prepared. The final delivered drug product will be mixed under aseptic conditions at a ratio of: 1 part (20%) of the active agents (in a concentration of between about .001 µg/ml to about 5 mg/ml) to 4 parts (80%) diluent.

10 The treatment regime will also vary depending on the disease being treated, based on a variety of factors, including the type of injury, the age, weight, sex, medical condition of the individual, the severity of the condition, the route of administration, and the particular compound employed. For example, the active agents are administered to an osteoporosis patient for up to 30 days. The therapy is administered  
15 for 1 to 6 times per day at dosages as described above.

In a preferred embodiment, the active agent is administered subcutaneously. A suitable subcutaneous dose of active ingredient of active agent is preferably between about 0.1 ng/kg and about 10 mg/kg administered twice daily for a time sufficient to enhance bone or cartilage repair. In a more preferred embodiment, the concentration of  
20 active agent is between about 100 ng/kg body weight and about 10.0 mg/kg body weight. In a most preferred embodiment, the concentration of active agent is between about 10 µg/kg body weight and about 10.0 mg/kg body weight. This dosage regimen maximizes the therapeutic benefits of the subject invention while minimizing the

amount of agonist needed. Such an application minimizes costs as well as possible deleterious side effects.

For subcutaneous administration, the active ingredient may comprise from 0.0001% to 10% w/w, *e.g.*, from 1% to 2% by weight of the formulation, although it  
5 may comprise as much as 10% w/w, but preferably not more than 5% w/w, and more preferably from 0.1% to 1% of the formulation.

In another preferred embodiment of the present invention, the active agent is administered topically at the site of bone or cartilage loss or repair. Suitable topical doses and active ingredient concentration in the formulation are as described for  
10 subcutaneous administration.

In another preferred embodiment, the active agent is administered at the desired site of bone or cartilage repair, such as via capsule delivery in gingival tissue at sites of bone resorption or in a scaffold surgically implanted at the site of a non-union bone fracture. Suitable doses and active ingredient concentration in the formulation are as  
15 described for subcutaneous administration.

In one embodiment of the invention, *ex vivo* methods are presented for enhancing bone repair via isolation of osteoblastic cell populations from a subject, contacting the isolated cell population with angiotensinogen, AI, AI analogues, AI fragments and analogues thereof, AII, AII analogues, AII fragments and analogues  
20 thereof and/or AII AT<sub>2</sub> type 2 receptor agonist, and subsequent reinfusion of the osteoblastic cell population into the subject. Methods for the isolation of osteoblastic cell population have been described. (Hiruma et al., 1997; Lamparter et al., 1998) In a preferred embodiment, human bone cells are cultured from outgrowths of trabecular bone fragments of the femoral head from patients undergoing hip replacement due to

fracture, and treated by several consecutive collagenase digestion periods, using 1 mg/ml type II collagenase solution (Worthington Diagnostic Systems) for 20 minutes per digestion period.

In another embodiment of the invention, *ex vivo* or *in vitro* methods are presented for enhancing cartilage repair via isolation of chondrocyte cell populations from a subject, contacting the isolated cell population with the active agent, and subsequent reinfusion of the chondrocyte cell population into the subject. Methods for the isolation of chondrocyte cell populations have been described. (Kato et al., J. Cell Biol., vol. 100, pages 477-485 (1985); U.S. Patent Nos. 4,642,120; 5,053,050; and 5,736,372, each herein incorporated by reference in its entirety).

In a preferred embodiment, autologous or homologous bone marrow is obtained by aspiration with a bone biopsy needle from the iliac crest or femoral canal. (U.S. Patent No. 5,053,050) The aspirated cells are injected into a phosphate buffered saline (PBS) containing 0.25% trypsin and injected sequentially through 17, 18 and 20 gauge needles to achieve a single cell suspension. The cells are plated in a density of 50-100 x 10<sup>6</sup> cells on 100 mm tissue culture dishes fed with BGJ b medium (GIBCO) with 15% F.C.S. (Fetal Calf Serum). The medium is changed daily or as required by the proliferation rate of the cells. The medium is supplemented with between about 0.1 ng/ml and 10 mg/ml active agent. The cells are subcultured weekly and after 5-6 subulturings an almost pure fibroblastic stromal cell population is achieved. This cell population is then trypsinized and put in a suspension culture at a density of 3-8 x 10<sup>6</sup> cells/ml of medium and cultured above soft agar in a F-12 medium with 10% F.C.S. and 50 µg/ml sodium ascorbate added daily to the medium. The fibroblastic stromal cells start to aggregate immediately and after three-seven days most of the cells are in

aggregates of 30-60 cells. All the aggregates express a chondrogenic phenotype, as determined by employing histochemical and immunohistochemical probes for analysis.

Although bone marrow derived chondrocytes are preferred, chondrocytes of autologous or homologous origin, or homologous committed chondrocytes, or any other progenital cells of mesenchymal origin can be used. It can be seen that this initial formulation comprises purification, proliferation and manipulation of a population expressing a chondrogenic phenotype. More specifically, the proliferating cells are from the class comprising bone marrow stroma cells, embryonal committed chondrocytes and any undifferentiated mesenchymal cells.

In a preferred embodiment, the proliferative effect of the active agents of the invention on cartilage cells is assessed by reactivity to an antibody directed against a protein known to be present in higher concentrations in proliferating cells than in non-proliferating cells, including but not limited to proliferating cell nuclear antigen (PCNA, or cyclin; Zymed Laboratories, South San Francisco, California). Viable cells may also be identified using a technique such as the trypan blue exclusion assay. Cells to be reinfused into the subject are rinsed to remove all traces of culture fluid, resuspended in an appropriate medium and then pelleted and rinsed several times. After the final rinse, the cells are resuspended at between  $0.7 \times 10^6$  and  $50 \times 10^6$  cells per ml in an appropriate medium and administered as described below.

Alternatively, the effects of the active agents on matrix component synthesis in bone and cartilage cells are determined in organ culture and in isolated cells by measuring the levels of two matrix proteins, as described in U.S. Patent No. 5,686,116 incorporated by reference herein in its entirety. Type I collagen is the major matrix protein of bone and cartilage. Collagen is almost entirely composed of proline, OH-

proline, alanine and glycine. Thus, new-bone collagen synthesis can be determined by measuring the uptake of  $^3\text{H}$ -Pro or by following the appearance of  $^3\text{H}$ -OH-Pro, which is formed by conversion of proline subsequent to its incorporation into collagen. Osteocalcin is a bone-specific matrix protein which is thought to be a cell signal for attracting osteoclasts to bone to initiate bone breakdown, and is also thought to slow or impede formation of newly mineralizing bone. Price et al., Proc. Natl. Acad. Sci. USA, 79:7734-7738 (1982). Therefore, decreased osteocalcin synthesis is associated with bone repair.

Any synoviocyte (cartilage-forming) cells can be used. In a preferred embodiment, the cell line HIG-82, a permanent cartilage cell line which retains many features present in normal synoviocytes, is used. Georgescu et al., In Vitro Cell. Dev. Biol., 24:1015-1022 (1988).

The synoviocytes are exposed to active agent for 24-48 hours, after which total cellular RNA is isolated by standard means. (*Molecular Cloning: A Laboratory Manual* (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press)) Detection of mRNA for type I collagen can be performed via standard techniques in the art including but not limited to reverse transcription-polymerase chain reaction (RT-PCR) using primers specific to type I collagen or osteocalcin, or Northern blotting of the RNA followed by hybridization with probes for type I collagen or osteocalcin. (*Molecular Cloning: A Laboratory Manual* (Sambrook, et al., 1989, Cold Spring Harbor Laboratory Press)).

Alternatively, the expression of type I collagen synthesis is determined in organ culture as described in U.S. Patent No. 5,686,116. In a preferred embodiment, calvarial (skull) bones from newborn rats are placed in sterile culture dishes with a nutritive

media to maintain viability. In this state, structural tissues grow by forming new matrix components (notably bone-specific collagen), but this growth is very slow. Kream et al., Endocrinol., 116:296-302 (1985). Hemicalvaria from 21 day old fetal rats are incubated for 48 hours in the presence and absence of angiotensinogen, AI, AI analogues, AI fragments and analogues thereof, AII, AII analogues, AII fragments and analogues thereof and/or AII AT<sub>2</sub> type 2 receptor agonist at various concentrations. <sup>3</sup>H-Pro is added for the final 18 hours of the incubation. Results showing that AII increases the levels of proline (Pro) and hydroxyproline (OH-Pro) in fetal rat hemicalvaria demonstrate that AII can increase the level of type I collagen synthesis in a normal bone culture system.

In another embodiment, the present invention comprises a method to enhance bone or cartilage repair *in vivo* by administration of the active agents of the invention. In one embodiment, the compounds of the invention are injected into the subcutaneous tissue over the right calvarium of mice. Control vehicle is PBS supplemented with 1% BSA. Heparin is administered at a dose of 50 units/ml. The animals are sacrificed on day 14 and bone growth is measured by histomorphometry.

Bone samples for quantitation are cleaned from adjacent tissues and fixed in 10% buffered formalin for 24-48 hours, decalcified in 14% EDTA for 1-3 weeks, processed through graded alcohols and embedded in paraffin wax. Three micron sections of the calvaria and femurs are prepared. Representative sections are selected for histomorphometric assessment of the effects of the active agent on bone formation and bone resorption. Sections are measured by using a camera attachment to directly trace the microscopic image onto a digitizing plate. Bone changes are measured on sections collected 200  $\mu$ m apart, over 4 adjacent 1 x 1 mm fields on both the injected



and noninjected sides of the calvaria. New bone is identified by its woven, rather than lamellar structure, and osteoclasts and osteoblasts are identified by their distinctive morphology. Histomorphometry software (Osteomeasure, Osteometrix, Inc., Atlanta) is used to process digitizer input to determine cell counts and feature areas or perimeters.

5           Alternatively, the active agents of the invention can be used to potentiate osteoblast function in intact animals. In a preferred embodiment, a model for abnormal osteoblast activity, weanling Sprague-Dawley rats, are placed on a phosphate and vitamin D-deficient diet as per the manufacturer's instructions (the diet, #80039, Teklad, Madison, Wis.) The animals on the diet are also kept in the dark to prevent de  
10       novo vitamin D synthesis. Animals placed under such conditions show abnormal bone formation and a marked deficiency in total bone mass.

          One group of the weanling rats on the diet is treated with active agent at between about 0.1 ng/kg and 10 mg/kg, given as a subcutaneous injection, every other day for 21 days. One group on the diet remains untreated and served as the control.  
15       Littermate controls not on the diet and not treated with active agent supply blood samples at the time of sacrifice of the animals on the diet for determination of alkaline phosphatase activity. Upon sacrifice, the long bones are removed from the animals on the diet for subsequent analyses.

          Serum alkaline phosphatase activity is used as a reliable indicator of osteoblast  
20       activity, such that increased levels of alkaline phosphatase activity are evidence of the abnormal bone turnover in the experimental animals. Serum alkaline phosphatase activity is determined by measuring the hydrolysis of p-nitrophenyl phosphate by serum samples according to the method of Lowry et al., J. Biol. Chem., 207:19-37 (1954). Markedly elevated serum alkaline phosphatase activity indicates abnormal

osteoblast activity. By contrast, normalization of bone cell function is evidenced by a decreased level of serum alkaline phosphate activity and increased bone mass (tested via determination of the ash weight of bones) relative to control animals.

Alternatively, a full thickness articular cartilage defect model in the femoral-  
5 patellar joint of adult rabbits is used to evaluate the ability of the compounds of the present invention to affect cartilage and bone repair, as described in U.S. Patent No. 5,700,774, hereby incorporated by reference in its entirety. Adult New Zealand White rabbits are anesthetized and prepared for sterile surgery. A 3 x 3 mm defect through  
10 articular cartilage and into underlying subchondral bone is drilled into the patellar groove of the knee joint. The defect is either left empty, filled with collagen sponge (controls), or with collagen sponge soaked with between about 0.1 ng/ml and 10 mg/ml active agent. The incision is closed and animals are allowed free movement within their cages for 4 weeks. After 4 weeks the animals are humanely euthanatized and the  
15 articular cartilage/subchondral bone defect is evaluated histologically for tissue architecture, quantity and quality of repair tissue. Northern analysis is performed for additional phenotyping.

In a further embodiment, dental and orthopedic implants can be coated with the compounds of the invention. In general, implant devices are coated with the active agents of the invention dissolved at a concentration in the range of 0.1 ng/ml to 10  
20 mg/ml in phosphate-buffered saline (PBS) containing 2 mg/ml serum albumin. The porous end of an implant is dipped in the solution and is air dried (or lyophilized) or implanted immediately into the bony site. The viscosity of the coating solution is increased, if desired, by adding hyaluronate at a final concentration of 0.1 mg/ml to 100 mg/ml or by adding other pharmaceutically acceptable excipients. Alternatively, the

solution containing the active agent is mixed with collagen gel or human collagen (e.g. Zyderm Registered TM Collagen Implant, Collagen Corp., Palo Alto, Calif.) to a final collagen concentration of 2 mg/ml to 100 mg/ml to form a paste or gel, which is then used to coat the porous end of the implant device. The coated implant device is placed  
5 into the bony site immediately or is air dried and rehydrated with PBS prior to implanting, with the objective of maximizing new bone formation into the implant while minimizing the ingrowth of soft tissue into the implant site.

In a further aspect, the present invention provides kits for enhancing bone or cartilage repair, wherein the kits comprise an effective amount of active agent for bone  
10 or cartilage repair, and instructions for using the amount effective of active agent as a therapeutic. In a preferred embodiment, the kit further comprises a pharmaceutically acceptable carrier, such as those adjuvants described above. In another preferred embodiment, the kit further comprises a means for delivery of the active agent to a patient. Such devices include, but are not limited to syringes, matrical or micellar  
15 solutions, bandages, wound dressings, polymeric scaffolds, collagen vehicles, aerosol sprays, lipid foams, transdermal patches, topical administrative agents, polyethylene glycol polymers, carboxymethyl cellulose preparations, crystalloid preparations (e.g., saline, Ringer's lactate solution, phosphate-buffered saline, etc.), viscoelastics, polyethylene glycols, and polypropylene glycols. The means for delivery may either  
20 contain the effective amount of the active agent, or may be separate from the compounds, which are then applied to the means for delivery at the time of use.

The kits may further comprise an amount effective for bone and/or cartilage repair, implantation, and augmentation of additional compounds, including but not limited to bone morphogenic protein-2, bone morphogenic protein-4, bone

morphogenic protein-6, bone morphogenic protein-7, transforming growth factor-beta, insulin-like growth factor, and parathyroid hormone. The kit may optionally contain a pharmaceutically acceptable carrier.

In another aspect of the present invention, an improved cell culture medium is provided for the proliferation of chondrocytes, wherein the improvement comprises addition to the cell culture medium of an effective amount of active agent to stimulate chondrocyte proliferation. Any cell culture media that can support the growth of chondrocytes can be used with the present invention. Such cell culture media include, but are not limited to Basal Media Eagle, Dulbecco's Modified Eagle Medium, Iscove's Modified Dulbecco's Medium, McCoy's Medium, Minimum Essential Medium, F-10 Nutrient Mixtures, Opti-MEM® Reduced-Serum Medium, RPMI Medium, and Macrophage-SFM Medium or combinations thereof.

The improved cell culture medium can be supplied in either a concentrated (ie: 10X) or non-concentrated form, and may be supplied as either a liquid, a powder, or a lyophilizate. The cell culture may be either chemically defined, or may contain a serum supplement. Culture media is commercially available from many sources, such as GIBCO BRL (Gaithersburg, MD) and Sigma (St. Louis, MO).

In another embodiment, the methods of the present invention can be used to treat chondrocytic cell lines, such as articular chondrocytes, in order to maintain chondrocytic phenotype and survival of the cells. The treated cell populations are therefore also useful for gene therapy applications.

In another preferred embodiment, the kit further comprises a sterile container. The sterile container can comprise either a sealed container, such as a cell culture flask,

a roller bottle, or a centrifuge tube, or a non-sealed container, such as a cell culture plate or microtiter plate (Nunc; Naperville, IL).

In a further preferred embodiment, the kit further comprises an antibiotic supplement for inclusion in the reconstituted cell growth medium. Examples of appropriate antibiotic supplements include, but are not limited to actinomycin D, Fungizone®, kanamycin, neomycin, nystatin, penicillin, streptomycin, or combinations thereof (GIBCO).

A further object of the present invention is to provide pharmaceutical compositions comprising the active agents as an ingredient for use in the methods of the invention. The compositions comprise the active agents together with a compound or compounds that also enhance bone or cartilage implantation, repair and regeneration, including, but not limited to bone morphogenic protein-2, bone morphogenic protein-4, bone morphogenic protein-6, bone morphogenic protein-7, transforming growth factor-beta, insulin-like growth factor, and parathyroid hormone, together with a pharmaceutically acceptable carrier, this term including any carrier which does not interfere with the effectiveness of the biological activity of the active agents and other compounds, and which is not toxic to the host to which it is administered. Dosage and administration of the pharmaceutical compositions will vary depending on the disease being treated, based on a variety of factors, including the type of injury, the age, weight, sex, medical condition of the individual, the severity of the condition, the route of administration, and the particular compound employed, as above. Thus, the dosage regimen may vary widely, but can be determined routinely by a physician using standard methods.

The present invention fulfills the need for methods to enhance bone and cartilage repair in a mammal suffering from a wide variety of disorders and injuries including, but not limited to bone fractures, defects, and disorders which result in weakened bones such as osteoporosis, osteoarthritis, Paget's disease, osteomalacia, periodontal disease; bone loss resulting from multiple myeloma and other forms of cancer; bone loss resulting from side effects of other medical treatment (such as steroids); age-related loss of bone mass; articular cartilage tears, deformities and other cartilage defects such as arthritis and cartilaginous tissue damage.

10 **Example 1. Method for Culture of Rabbit Chondrocytes**

Chondrocytes were isolated from the cartilage of the knee joints of adult rabbits and cultured according to the method of Okazaki et al. (Ann. Rheum. Dis. 55:181-186, 1996). Briefly, cartilage explants were minced into small pieces (approximately 1.5 mm by 1.5 mm). The tissue pieces were digested at 37°C sequentially in 0.05% hyaluronidase (415 U/mg protein) for 10 minutes, 0.2% Type III trypsin (10,000 U/mg protein, Sigma) and 0.53 mM EDTA for 15 minutes, and 0.2% Type I collagenase (136 U/mg protein, Sigma) for 30 minutes. The samples were then washed and incubated overnight at 37°C in 5% CO<sub>2</sub> in air in Dulbecco's modified Eagle's medium enriched with 10% fetal calf serum, 3.5 mg/ml glucose, 0.2% Type I collagenase and antibiotics (100 U/ml penicillin and 100 µg/ml streptomycin).

After this incubation, the cells were harvested and washed once with phosphate buffered saline (pH 7.2). The cells were then counted with a hemacytometer and resuspended at  $1 \times 10^5$  cells/ml in Ham's F12 supplemented with 10% fetal calf serum and antibiotics. A 10 ml aliquot of cells was transferred to collagen coated 25 cm<sup>2</sup>

flasks (coated with collagen isolated from rat tail tendons) and incubated at 37°C in 5% CO<sub>2</sub> in air. Five to seven days after the initiation of culture, the cells were detached with 0.05% trypsin-EDTA (Gibco-BRL) at 37°C in 5% CO<sub>2</sub> in air. After detachment, the cells were washed once with phosphate buffered saline, centrifuged and resuspended at 200 cells/ml in Ham's F12 supplemented with 10% fetal calf serum and antibiotics. Two hundred microliters of cells were then aliquoted into the collagen coated wells of a 96 well microtiter plate and allowed to adhere, after which, 10 µg/ml of the AII and AII analogue and fragment peptides listed in Table 3 were added to the wells to assess their effects on chondrocyte proliferation on days 1, 2, and 3 after culture initiation. Cell numbers were quantitated by staining the cells with Giemsa stain and counting the number of nuclei detected via microscopy. The data (FIG. 1-4) demonstrate that AII and the AII analogues and fragments all accelerated the chondrocyte proliferation.

**Table 3. Designation for Analogues/Fragments Used in Example 1**

Name	Abbreviation	Sequence	SEQ ID NO:
GSD 37B	Orn <sup>2</sup> -AII	D(Orn)VYIHPF	SEQ ID NO:38
GSD 28	Ile <sup>8</sup> -AII	DRVYIHPI	SEQ ID NO:39
GSD 24B	Pro <sup>3</sup> -AII	DRPYIHPF	SEQ ID NO:31
GSD 22A	Ala <sup>4</sup> -AIII	RVYAHPF	SEQ ID NO:18
GSD36	Gly <sup>1</sup> -AII	GRVYIHPF	SEQ ID NO:42
GSD38B	Citron <sup>2</sup> -AII	D(Citron)VYIHPF	SEQ ID NO:43
1GD	Ala <sup>4</sup> -AII(1-7)	DRVAIHP	SEQ ID NO:40
2GD	Pro <sup>3</sup> -AII(1-7)	DRPYIHP	SEQ ID NO:41
3GD	Pro <sup>3</sup> Ala <sup>4</sup> -AII(1-7)	DRPAIHP	SEQ ID NO:44
AIII	AII(2-8)	RVYIHPF	SEQ ID NO:2
AII(1-7)		DRVYIHP	SEQ ID NO:4
AII		DRVYIHPF	SEQ ID NO. 1

**Example 2. Bone Healing**

Female, Sprague Dawley rats underwent intramuscular anesthesia with ketamine/rompum and were prepared for sterile surgery by shaving the surgical site and scrubbing with Betadine scrub followed by 70% ethanol. The rat was then placed on a sterile field in a lateral decubitus position facing the surgeon. The shaved legs were then covered with Betadine solution and draped aseptically. A skin incision was performed parallel to the long axis of the right medial diaphysis. The muscle was separated along fascial planes to expose the tibia. A defect of 1.4 mm in diameter was then drilled from the lateral side of the midshaft cortex so that the defect extended from one cortical side to the other, through the bone marrow. Sterile saline (0.9% NaCl) for injection was then used to clean the surgical area of tissue debris and bone fragments. Either hydron polymer solution (vehicle: 10% Hydron, 60% ethanol, 1% polyethylene glycol polymer) or peptide (AII, AII(1-7), or 9GD at 1 mg/ml) in hydron polymer solution was placed in the bone defect to fill the defect with polymer (approximately 0.1 ml of polymer). The incision was closed with 3-0 Vicryl suture using continuous mattress suture. The animals were allowed to recover from anesthesia, given Bupronex for analgesia and allowed free movement, until euthanasia 7 days later.

By gross observation, the defects that received the peptides of the invention were more completely filled with new tissue that had begun to calcify. The majority of control defects were less than 50% filled with new tissue and no hardening of the tissue was observed. These data demonstrate that the peptides of the invention accelerate the formation of new bone tissue.



After gross evaluation, the muscle tissue was removed from the bone and the bones were placed in formalin for fixation. After 2 days in 10% buffered formaldehyde, the tissues were placed in a decalcifying solution (Rapid Bone Decal, American MasterTech Scientific, Inc. Lodi,

5 CA) diluted 3:1 for 6 hours. After decalcification, the bone was cut in half along the long axis, embedded in paraffin, sectioned and stained with hematoxylin and eosin.

Microscopic evaluation of the tissue sections confirmed the gross observations. On day 7, the bones in which the defect had been filled with Hydron (the placebo) had a loose fibrin filler with the majority of cells observed being inflammatory in nature.

10 Occasionally, a cell that appeared to be of mesenchymal origin or a blood vessels was seen at the site of injury. In all of the peptide treated animals, extensive stromal cell ingrowth with numerous blood vessels was observed. In approximately 50% of the peptide treated animals, tissue with the appearance of new bone was observed. These data clearly support the ability of these angiotensin peptides to accelerate new bone  
15 formation.

**Table 4. Designation for Analogues/Fragments used in Example 2**

20	AII(1-7)	DRVYIHP	SEQ ID NO:4
	AII	DRVYIHPF	SEQ ID NO. 1
	9GD:	NorLeu3-AII(1-7) DR(nor)YIHP	SEQ ID NO:45

It is to be understood that the invention is not to be limited to the exact details of operation, or to the exact compounds, compositions, methods, procedures or  
25 embodiments shown and described, as obvious modifications and equivalents will be

apparent to one skilled in the art, and the invention is therefore to be limited only by the full scope of the appended claims.

We claim:

1. A method for enhancing bone repair in a mammal, comprising the administration of an amount effective for enhancing bone repair in the mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent.

2. An improved method for bone and prosthesis implantation in a mammal, comprising the administration of an amount effective for enhancing bone and prosthesis implantation in the mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

15  $R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

20

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent.

3. A kit for enhancing bone repair in a mammal, comprising:

- (a) administering an amount effective for enhancing bone repair in the mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



in which  $R^1$  and  $R^2$  together form a group of formula



10 wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

20  $R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;


$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent; and

(b) instructions for using the amount effective of active agent to enhance bone repair in a mammal.

5 4. A kit for improved bone and prosthesis implantation, comprising:

(a) an amount effective to enhance bone and prosthesis implantation in a mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



10 in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

20  $R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent; and

- (b) instructions for using the amount effective of active agent to enhance  
5 bone and prosthesis implantation in a mammal.

5. A method for enhancing cartilage repair in a mammal, comprising the administration of an amount effective to enhance cartilage repair to the mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I

10  $R^1$ - $R^2$ - $R^3$ - $R^4$ - $R^5$ - $R^6$ - $R^7$ - $R^8$ .

in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly,  
15 Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile,  
20 Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent.

5 6. A kit for enhancing cartilage repair in a mammal, comprising:

(a) an amount effective to enhance cartilage repair in a mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



10 in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

20  $R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;



$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent; and

(b) instructions for using the amount effective of active agent for enhancing  
5 cartilage repair in a mammal.

7. An improved method for attachment and fixation of cartilage implants to bone or other tissue in a mammal, comprising the administration of an amount effective for attaching and fixing cartilage implants to bone or other tissue in a mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino  
10 acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

15  $R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

20  $R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;


$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

5  $R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent.

8. A kit for improved attachment and fixation of cartilage implants to bone or other tissue, comprising:

(a) an amount effective to attach and fix a cartilage implant to bone or other  
10 tissue in a mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



in which  $R^1$  and  $R^2$  together form a group of formula



15 wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;  
20

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;


$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

5  $R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent; and

(b) instructions for using the amount effective of active agent amount effective to attach and fix a cartilage implant to bone or other tissue in a mammal.

9. An improved medium for the culture of chondrocytes wherein the improvement  
10 comprises contacting the cells with an amount effective to accelerate proliferation of chondrocytes of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

20  $R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

5  $R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent.

10. The improved medium of claim 9 wherein the active agent is selected from the  
 10 group consisting of angiotensinogen, SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID  
 15 NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO: 34; SEQ ID NO:35, SEQ ID NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:45.

11. A kit for the culture of chondrocytes, comprising

20 (a) an amount effective to accelerate proliferation of chondrocytes of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I



in which  $R^1$  and  $R^2$  together form a group of formula



wherein X is H or a one to three peptide group, or is absent

$R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

$R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent; and

(b) instructions for using the amount effective of active agent amount effective to accelerate proliferation of chondrocytes.

12. The method of claim 1, 2, 5, or 7 wherein the active agent is selected from the group consisting of angiotensinogen, SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID

NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID  
NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID  
NO:31, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO: 34; SEQ ID NO:35, SEQ ID  
NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID  
5 NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:45.

13. The kit of claim 3, 4, 6, 8, or 11 wherein the active agent is selected from the  
group consisting of angiotensinogen, SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:3,  
SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:8, SEQ ID  
NO:9, SEQ ID NO:10, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:13, SEQ ID  
10 NO:16, SEQ ID NO:17, SEQ ID NO:18, SEQ ID NO:19, SEQ ID NO:20, SEQ ID  
NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID  
NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID  
NO:31, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO: 34; SEQ ID NO:35, SEQ ID  
NO:36, SEQ ID NO:37, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID  
15 NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:45.

14. A method for enhancing bone repair in a mammal, comprising the  
administration of an amount effective for enhancing bone repair in the mammal of at  
least one active agent of the formula

Asp-Arg-R1-R2-Ile-His-Pro-R2, wherein

20 R1 is selected from the group consisting of Ile, Pro, Ala, Val, Leu, and norLeu;

R2 is selected from Tyr and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe, or is absent.

15. An improved method for bone and prosthesis implantation in a mammal, comprising the administration of an amount effective for enhancing bone and prosthesis implantation to the mammal of at least one active agent of the formula

Asp-Arg-R1-R2-Ile-His-Pro-R2, wherein

5 R1 is selected from the group consisting of Ile, Pro, Ala, Val, Leu, and norLeu;

R2 is selected from Tyr and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe, or is absent.

16. The method of claim 15 or 16 wherein the active agent is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:24, SEQ ID NO:31,

10 SEQ ID NO:32, SEQ ID NO: 33, SEQ ID NO:41, and SEQ ID NO:45.

17. A kit for enhancing bone repair in a mammal, comprising:

(a) an amount effective for enhancing bone repair in the mammal of at least one active agent of the formula

Asp-Arg-R1-R2-Ile-His-Pro-R2, wherein

15 R1 is selected from the group consisting of Ile, Pro, Ala, Val, Leu, and norLeu;

R2 is selected from Tyr and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe, or is absent; and

(b) instructions for using the amount effective of active agent to enhance bone repair in a mammal.

20

18. A kit for improved bone and prosthesis implantation, comprising:

(a) an amount effective to enhance bone and prosthesis in a mammal of at least one active agent of the formula

Asp-Arg-R1-R2-Ile-His-Pro-R2, wherein

R1 is selected from the group consisting of Ile, Pro, Ala, Val, Leu, and norLeu;

R2 is selected from Tyr and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe, or is absent; and

- (b) instructions for using the amount effective of active agent to enhance  
5 bone and prosthesis implantation in a mammal.
19. The kit of claim 17 or 18 wherein the active agent is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:4, SEQ ID NO:24, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO: 33, SEQ ID NO:41, and SEQ ID NO:45.
- 10 20. A method for enhancing cartilage repair in a mammal, comprising the administration of an amount effective to enhance cartilage repair to the mammal of at least one active agent of the formula

R1-R2-R3-R4-R5-His-Pro-R6, wherein

R1 is selected from the group consisting of Hydrogen, Gly, and Asp;

15 R2 is selected from the group consisting of Arg, Citron, or Ornithine;

R3 is selected from the group consisting of Val, Ile, Ala, Leu, and norLeu, or Pro;

R4 is selected from Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, and Ala;

R5 is selected from the group consisting of Ile, Ala, Val, Leu, and norLeu; and

20 R6 is Phe, Ile, or is absent.

21. A kit for enhancing cartilage repair in a mammal, comprising:

- (a) an amount effective to enhance cartilage repair in a mammal of at least one active agent comprising a sequence of the formula



R1-R2-R3-R4-R5-His-Pro-R6, wherein

R1 is selected from the group consisting of Hydrogen, Gly, and Asp;

R2 is selected from the group consisting of Arg, Citron, or Ornithine;

R3 is selected from the group consisting of Val, Ile, Ala, Leu, and norLeu, or

5 Pro;

R4 is selected from Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, and Ala;

R5 is selected from the group consisting of Ile, Ala, Val, Leu, and norLeu; and

R6 is Phe, Ile, or is absent; and

(b) instructions for using the amount effective of active agent for enhancing  
10 cartilage repair in a mammal.

22. An improved method for attachment and fixation of cartilage implants to bone  
or other tissue in a mammal, comprising the administration of an amount effective for  
attaching and fixing cartilage implants to bone or other tissue in a mammal of at least  
15 one active agent of the formula

R1-R2-R3-R4-R5-His-Pro-R6, wherein

R1 is selected from the group consisting of Hydrogen, Gly, and Asp;

R2 is selected from the group consisting of Arg, Citron, or Ornithine;

R3 is selected from the group consisting of Val, Ile, Ala, Leu, and norLeu, or

20 Pro;

R4 is selected from Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, and Ala;

R5 is selected from the group consisting of Ile, Ala, Val, Leu, and norLeu; and

R6 is Phe, Ile, or is absent.

23. A kit for improved attachment and fixation of cartilage implants to bone or other tissue, comprising:

(a) an amount effective to attach and fix a cartilage implant to bone or other tissue in a mammal of at least one active agent comprising a sequence of the formula

5 R1-R2-R3-R4-R5-His-Pro-R6, wherein

R1 is selected from the group consisting of Hydrogen, Gly, and Asp;

R2 is selected from the group consisting of Arg, Citron, or Ornithine;

R3 is selected from the group consisting of Val, Ile, Ala, Leu, and norLeu, or Pro;

10 R4 is selected from Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, and Ala;

R5 is selected from the group consisting of Ile, Ala, Val, Leu, and norLeu; and

R6 is Phe, Ile, or is absent; and

(b) instructions for using the amount effective of active agent amount  
15 effective to attach and fix a cartilage implant to bone or other tissue in a mammal.

24. The method of claim 20 or 22 wherein the active agent is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 13, SEQ ID NO: 18, SEQ ID NO:19, SEQ ID NO:12, SEQ ID NO:24, SEQ ID NO:26,  
20 SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:45.

25. An improved medium for the culture of chondrocytes wherein the improvement comprises contacting the cells with an amount effective to accelerate proliferation of chondrocytes of at least one active agent comprising a sequence of the formula

R1-R2-R3-R4-R5-His-Pro-R6, wherein

5 R1 is selected from the group consisting of Hydrogen, Gly, and Asp;

R2 is selected from the group consisting of Arg, Citron, or Ornithine;

R3 is selected from the group consisting of Val, Ile, Ala, Leu, and norLeu, or Pro;

R4 is selected from Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, and Ala;

10 R5 is selected from the group consisting of Ile, Ala, Val, Leu, and norLeu; and

R6 is Phe, Ile, or is absent.

26. The medium of claim 25 wherein the active agent is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 13, SEQ ID NO: 18, SEQ ID NO:19, SEQ ID NO:12, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:45.

27. A kit for the culture of chondrocytes, comprising

(a) an amount effective to accelerate proliferation of chondrocytes of at least one active agent comprising a sequence of the formula

R1-R2-R3-R4-R5-His-Pro-R6, wherein

R1 is selected from the group consisting of Hydrogen, Gly, and Asp;

R2 is selected from the group consisting of Arg, Citron, or Ornithine;

R3 is selected from the group consisting of Val, Ile, Ala, Leu, and norLeu, or Pro;

R4 is selected from Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, and Ala;

R5 is selected from the group consisting of Ile, Ala, Val, Leu, and norLeu; and

5 R6 is Phe, Ile, or is absent; and

(b) instructions for using the amount effective of active agent amount effective to accelerate proliferation of chondrocytes.

28. The kit of claim 21, 23, or 27 wherein the active agent is selected from the  
10 group consisting of SEQ ID NO:1, SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO: 13, SEQ ID NO: 18, SEQ ID NO:19, SEQ ID NO:12, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO: 32, SEQ ID NO:33, SEQ ID NO:38, SEQ ID NO:39, SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42, SEQ ID NO:43, SEQ ID NO:44, and SEQ ID NO:45.

15 29. The method of claim 1, 2, 5, 7, 12, 14, 15, 16, 20, 22, or 24 wherein the concentration of active agent is between about 0.1 ng/kg and about 1.0 mg/kg.

30. The kit of claim 3, 4, 6, 8, 13, 17, 18, 19, 21, 23, or 28 wherein the concentration of active agent is between about 0.1 ng/kg and about 1.0 mg/kg.

31. The kit of claim 11 or 27 wherein the concentration of active agent is between  
20 about 0.1 ng/ml and about 1.0 mg/ml.

32. The kit of claim 11 or 27 further comprising tissue culture medium.

33. The kit of claim 3, 4, 6, 8, 13, 17, 18, 19, 21, 23, or 28 further comprising a means for delivery of the active agent.

34. The culture medium of claim 9, 10, 25, or 26 wherein the concentration of active agent is between about 0.1 ng/ml and about 1.0 mg/ml.

35. The kit of claim 3, 4, 6, 8, 13, 17, 18, 19, 21, 23, or 28 further comprising an amount effective for enhancing bone or cartilage repair, regeneration, or implantation in the mammal of at least one compound selected from bone morphogenic protein-2, bone morphogenic protein-4, bone morphogenic protein-6, bone morphogenic protein-7, transforming growth factor-beta, insulin-like growth factor, and parathyroid hormone.

36. The method of claim 1, 2, 5, 7, 12, 14, 15, 16, 20, 22, or 24 further comprising administering an amount effective for enhancing bone or cartilage repair, regeneration, or implantation in the mammal of at least one compound selected from bone morphogenic protein-2, bone morphogenic protein-4, bone morphogenic protein-6, bone morphogenic protein-7, transforming growth factor-beta, insulin-like growth factor, and parathyroid hormone.

37. A pharmaceutical composition comprising

(a) an amount effective to enhance bone or cartilage repair, bone and prosthesis implantation, or cartilage implantation in a mammal of at least one active agent comprising a sequence of the formula

Asp-Arg-R1-R2-Ile-His-Pro-R2, wherein

R1 is selected from the group consisting of Ile, Pro, Ala, Val, Leu, and norLeu;

R2 is selected from Tyr and Tyr(PO<sub>3</sub>)<sub>2</sub>; and

R3 is Phe, or is absent;

(b) an amount effective to enhance bone repair or bone and prosthesis implantation in a mammal of at least one compound selected from bone morphogenic

protein-2, bone morphogenic protein-4, bone morphogenic protein-6, bone morphogenic protein-7, transforming growth factor-beta, insulin-like growth factor, and parathyroid hormone; and

(c) a pharmaceutically acceptable carrier.

5 38. A pharmaceutical composition comprising

(a) an amount effective to enhance bone repair or bone and prosthesis implantation in a mammal of at least one active agent comprising a sequence consisting of at least three contiguous amino acids of groups  $R^1$ - $R^8$  in the sequence of general formula I

10  $R^1$ - $R^2$ - $R^3$ - $R^4$ - $R^5$ - $R^6$ - $R^7$ - $R^8$

in which  $R^1$  and  $R^2$  together form a group of formula

$X$ - $R^A$ - $R^B$ ,

wherein  $X$  is H or a one to three peptide group, or is absent

15  $R^A$  is suitably selected from H, Asp, Glu, Asn, Acpc (1-aminocyclopentane carboxylic acid), Ala, Me<sup>2</sup>Gly, Pro, Bet, Glu(NH<sub>2</sub>), Gly, Asp(NH<sub>2</sub>) and Suc,

$R^B$  is suitably selected from Arg, Lys, Ala, Orn, Citron, Ser(Ac), Sar, D-Arg and D-Lys;

20  $R^3$  is selected from the group consisting of Val, Ala, Leu, norLeu, Ile, Gly, Pro, Aib, Acpc and Tyr;

$R^4$  is selected from the group consisting of Tyr, Tyr(PO<sub>3</sub>)<sub>2</sub>, Thr, Ala, Ser, homoSer and azaTyr;

$R^5$  is selected from the group consisting of Ile, Ala, Leu, norLeu, Val and Gly;

$R^6$  is His, Arg or 6-NH<sub>2</sub>-Phe;

$R^7$  is Pro or Ala; and

$R^8$  is selected from the group consisting of Phe, Phe(Br), Ile and Tyr, excluding sequences including  $R^4$  as a terminal Tyr group, or is absent;

- 5           (b)    an amount effective to enhance bone repair or bone and prosthesis implantation in a mammal of at least one compound selected from bone morphogenic protein-2, bone morphogenic protein-4, bone morphogenic protein-6, bone morphogenic protein-7, transforming growth factor-beta, insulin-like growth factor, and parathyroid hormone; and
- 10           (c)    a pharmaceutically acceptable carrier.

1/4




FIG 1



2/4




FIG 2

3/4




FIG 3

4/4




FIG 4

## SEQUENCE LISTING

<110> Rodgers, Kathleen  
diZerega, Gere

<120> Methods for Accelerating Bone and Connective Tissue  
Growth and Repair

<130> 98365C

<140> To be assigned

<141> 1999-07-11

<160> 45

<170> PatentIn Ver. 2.0

<210> 1

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII

<400> 1

Asp Arg Val Tyr Ile His Pro Phe

1

5

<210> 2

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (2-8)

<400> 2

Arg Val Tyr Ile His Pro Phe

1

5

<210> 3

<211> 6

<212> PRT

<213> Artificial Sequence

<223> Description of Artificial Sequence: AII (3-8)

<400> 3

Val Tyr Ile His Pro Phe  
1 5

<210> 4

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (1-7)

<400> 4

Asp Arg Val Tyr Ile His Pro  
1 5

<210> 5

<211> 6

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (2-7)

<400> 5

Arg Val Tyr Ile His Pro  
1 5

<210> 6

<211> 5

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII (3-7)

<400> 6

Val Tyr Ile His Pro  
1 5

<210> 7

<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (5-8)

<400> 7  
Ile His Pro Phe  
1

<210> 8  
<211> 6  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (1-6)

<400> 8  
Asp Arg Val Tyr Ile His  
1 5

<210> 9  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (1-5)

<400> 9  
Asp Arg Val Tyr Ile  
1 5

<210> 10  
<211> 4  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (1-4)

<400> 10  
Asp Arg Val Tyr

<210> 11  
<211> 3  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (1-3)

<400> 11  
Asp Arg Val  
1

<210> 12  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue

<220>  
<221> MOD\_RES  
<222> (2)  
<223> Nle

<400> 12  
Arg Xaa Tyr Ile His Pro Phe  
1 5

<210> 13  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue

<220>  
<221> MOD\_RES  
<222> (4)  
<223> Nle

<400> 13

Arg Val Tyr Xaa His Pro Phe  
1 5

<210> 14  
<211> 3  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (6-8)

<400> 14  
His Pro Phe  
1

<210> 15  
<211> 5  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII (4-8)

<400> 15  
Tyr Ile His Pro Phe  
1 5

<210> 16  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue  
class

<220>  
<221> UNSURE  
<222> (1)  
<223> Xaa at position 1 can be Arg, Lys, Ala, Orn, Ser,  
MeGly, D-Arg, or D-Lys

<220>  
<221> UNSURE  
<222> (2)



<223> Xaa at position 2 can be Val, Ala, Leu, Nle, Ile,  
Gly, Pro, Aib, Acp, or Tyr

<220>

<221> UNSURE

<222> (4)

<223> Xaa at position 4 can be Ile, Ala, Leu, Nle, Val,  
or Gly

<400> 16

Xaa Xaa Tyr Xaa His Pro Phe

1

5

<210> 17

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue

<400> 17

Arg Val Tyr Gly His Pro Phe

1

5

<210> 18

<211> 7

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue

<400> 18

Arg Val Tyr Ala His Pro Phe

1

5

<210> 19

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 1

Asp Arg Val Tyr Val His Pro Phe  
1 5

<210> 20  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 2

<400> 20  
Asn Arg Val Tyr Val His Pro Phe  
1 5

<210> 21  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 3

<400> 21  
Ala Pro Gly Asp Arg Ile Tyr Val His Pro Phe  
1 5 10

<210> 22  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 4

<400> 22  
Glu Arg Val Tyr Ile His Pro Phe  
1 5

<210> 23  
<211> 8  
<212> PRT  
<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: AII analogue 5

&lt;400&gt; 23

Asp Lys Val Tyr Ile His Pro Phe

1

5

&lt;210&gt; 24

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: AII analogue 6

&lt;400&gt; 24

Asp Arg Ala Tyr Ile His Pro Phe

1

5

&lt;210&gt; 25

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: AII analogue 7

&lt;400&gt; 25

Asp Arg Val Thr Ile His Pro Phe

1

5

&lt;210&gt; 26

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence: AII analogue 8

&lt;400&gt; 26

Asp Arg Val Tyr Leu His Pro Phe

1

5

<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 9

<400> 27  
Asp Arg Val Tyr Ile Arg Pro Phe  
1 5

<210> 28  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 10

<400> 28  
Asp Arg Val Tyr Ile His Ala Phe  
1 5

<210> 29  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 11

<400> 29  
Asp Arg Val Tyr Ile His Pro Tyr  
1 5

<210> 30  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 12

<400> 30

Pro Arg Val Tyr Ile His Pro Phe  
1 5

<210> 31

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 13

<400> 31

Asp Arg Pro Tyr Ile His Pro Phe  
1 5

<210> 32

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 14

<220>

<221> MOD\_RES

<222> (4)

<223> PHOSPHORYLATION

<400> 32

Asp Arg Val Tyr Ile His Pro Phe  
1 5

<210> 33

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: AII analogue 15

<220>

<221> MOD\_RES

<222> (3)

<223> Nle

Asp Arg Xaa Tyr Ile His Pro Phe  
1 5

<210> 34  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 16

<220>  
<221> MOD\_RES  
<222> (5)  
<223> Nle

<400> 34  
Asp Arg Val Tyr Xaa His Pro Phe  
1 5

<210> 35  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence: AII analogue 17

<220>  
<221> MOD\_RES  
<222> (4)  
<223> homo Ser

<400> 35  
Asp Arg Val Ser Tyr Ile His Pro Phe  
1 5

<210> 36  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial

Sequence:p-aminophenylalanine 6 AII

<220>

<221> MOD\_RES

<222> (6)

<223> p-aminophenylalanine

<400> 36

Asp Arg Val Tyr Ile Xaa Pro Phe

1

5

<210> 37

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:angiotensin I

<400> 37

Asp Arg Val Tyr Ile His Pro Phe His Leu

1

5

10

<210> 38

<211> 8

<212> PRT

<213> Artificial Sequence

<220>

<221> MOD\_RES

<222> (2)

<223> Orn

<220>

<223> Description of Artificial Sequence:GSD37B:  
Orn2-AII

<400> 38

Asp Xaa Val Tyr Ile His Pro Phe

1

5

<210> 39

<211> 8

<212> PRT

<213> Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:GSD28: Ile8-AII

&lt;400&gt; 39

Asp Arg Val Tyr Ile His Pro Ile

1

5

&lt;210&gt; 40

&lt;211&gt; 7

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:1GD:

Ala4-AII(1-7)

&lt;400&gt; 40

Asp Arg Val Ala Ile His Pro

1

5

&lt;210&gt; 41

&lt;211&gt; 7

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:2GD:

Pro3-AII(1-7)

&lt;400&gt; 41

Asp Arg Pro Tyr Ile His Pro

1

5

&lt;210&gt; 42

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; Description of Artificial Sequence:Gly1-AII

&lt;400&gt; 42

Gly Arg Val Tyr Ile His Pro Phe

1

5



<210> 43  
<211> 8  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial  
Sequence:GSD38B:Citron2-AII

<220>  
<221> MOD\_RES  
<222> (2)  
<223> Citron

<400> 43  
Asp Xaa Val Tyr Ile His Pro Phe  
1 5

<210> 44  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial  
Sequence:Pro3Ala4-AII(1-7)

<400> 44  
Asp Arg Pro Ala Ile His Pro  
1 5

<210> 45  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220>  
<223> Description of Artificial Sequence:9GD:  
norleu3-AII(1-7)

<220>  
<221> MOD\_RES  
<222> (3)  
<223> Nle

<400> 45

Asp Arg Xaa Tyr Ile His Pro

1

5